

Synthesis and nonlinear optical properties of pyridoxine-based stilbazole derivatives and their azo-analogs

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018, Copyright © Kazan (Volga region) Federal University. The syntheses of a series of novel derivatives of stilbazole and their azo-analogs bearing the phenylalkenyl and phenyldiazenyl substituents at position 6 of the pyridoxine ring are reported. The observed cis/trans regioselectivity in the Wittig reaction between triphenylbenzylphosphonium chloride and carbonyl derivatives of pyridoxine ketal is discussed with respect to transition state intermediates, nature of base used in the reaction, and resonance transformations in the pyridoxine system. The azo-derivatives are obtained by azocoupling of the diazonium salts of sulfanilic or 4-aminophenyl-1,3-disulfonic acids with the corresponding pyridoxine acetals. Nine compounds from the obtained experimental series are capable of emitting second harmonic generation light, although with a reduced efficiency as compared to well known KH_2PO_4 or LiIO_3 perovskite crystals, but with excellent beam resistance under laser irradiation, which makes them potential starting points for the development of novel nonlinear optical materials.

<http://dx.doi.org/10.1080/00397911.2017.1422520>

Keywords

Azo-derivatives of pyridoxine, derivatives of stilbazole, second harmonic generation

References

- [1] Verbiest, T.,; Houbrechts, S.,; Kauranen, M.,; Clays, K.,; Persoons, A., Second-Order Nonlinear Optical Materials: Recent Advances in Chromophore Design. *Phys. Rev. Lett.* 1997, 7, 2175– 2189. DOI: 10.1039/a703434b.
- [2] Cox, G., Biological Applications of Second Harmonic Imaging. *Biophys. Rev.* 2011, 3, 131– 141. DOI: 10.1007/s12551-011-0052-9.
- [3] Chung, I.,; Kanatzidis, G. M., Metal Chalcogenides: A Rich Source of Nonlinear Optical Materials. *Chem. Mater.* 2014, 26, 849– 869. DOI: 10.1021/cm401737s.
- [4] Nath, G.,; Haussuhl, S., Large Nonlinear Optical Coefficient and Phase- Matched Second Harmonic Generation in LiIO_3 . *Appl. Phys. Lett.* 1969, 14, 154– 156. DOI: 10.1063/1.1652754.
- [5] Suresh, S.,; Ramanand, A.,; Jayaraman, D.,; Mani, P., Review on Theoretical Aspect of Nonlinear Optics. *Rev. Adv. Mater. Sci.* 2012, 30, 175– 183.
- [6] (a) Wang, A.,; Yu, W.,; Huang, Z.,; Zhou, F.,; Song, J.,; Song, Y.,; Long, L.,; Cifuentes, M. P.,; Humphrey, M. G.,; Zhang, L.,; et al. Covalent Functionalization of Reduced Graphene Oxide with Porphyrin by Means of Diazonium Chemistry for Nonlinear Optical Performance. *Sci. Rep.* 2016, 6, 1– 12. DOI: 10.1038/srep23325;
- [7] (b) Zhao, W.,; Wang, A.,; Wang, Y.,; Lv, C.,; Zhu, W.,; Dou, S.,; Wang, Q.,; Zhong, Q., Accessible Fabrication and Mechanism Insight of Heterostructured $\text{BiOCl}/\text{BiMoO}_6/\text{g-CN}$ Nanocomposites with Efficient Photosensitized Activity. *J. Alloys Compd.* 2017, 726, 164– 172. DOI: 10.1016/j.jallcom.2017.07.268;

- [8] (c) Zhao, W.; Wang, Y.; Wang, A.; Qian, J.; Zhu, W.; Dou, S.; Wang, Q.; Zhong, Q.; Chen, A., Novel BiOCO/polypyrrole/g-CN Nanocomposites With Efficient Photocatalytic and Nonlinear Optical Properties. *RSC Adv.* 2017, 7, 7658– 7670.
- [9] (a) Le Bozec, H.; Renouard, T., Dipolar and Non-Dipolar Pyridine and Bipyridine Metal Complexes for Nonlinear Optics. *Eur. J. Inorg. Chem.* 2000, 2000, 229– 239. DOI: 10.1002/(sici)1099-0682(200002)2000:2<229::ai-ejic229>3.0.co;2-a;
- [10] (b) Cariati, E.; Pizzotti, M.; Roberto, D.; Tessore F.; Ugo, R., Coordination and Organometallic Compounds and Inorganic-Organic Hybrid Crystalline Materials for Second-Order Non-Linear Optics. *Coord. Chem. Rev.* 2006, 250, 1210– 1233. DOI: 10.1016/j.ccr.2005.09.013;
- [11] (c) Dragonetti, C.; Valore, A.; Colombo, A.; Righetto, S.; Rampinini, G.; Colombo, F.; Rocchigiani, L.; Macchioni, A., An Investigation on the Second-Order NLO Properties of Novel Cationic Cyclometallated Ir(III) Complexes of the Type [Ir(2-phenylpyridine)(9-R-4,5-diazafluorene)] (R⁺H, fulleridene) and the Related Neutral Complex with the New 9-Fulleriden-4-Monoazafluorene Ligand. *Inorganica Chim. Acta.* 2012, 382, 72– 78. DOI: 10.1016/j.ica.2011.10.018;
- [12] (d) Colombo, A.; Locatelli, D.; Roberto, D.; Tessore, F.; Ugo Righetto, S.; Rampinini, G.; Colombo, F.; Rocchigiani, L.; Macchioni, A., An Investigation on the Second-Order NLO Properties of Novel Cationic Cyclometallated Ir(III) Complexes of the Type [Ir(2-phenylpyridine)(9-R-4,5-diazafluorene)] (R⁺H, fulleridene) and the Related Neutral Complex with the New 9-Fulleriden-4-Monoazafluorene Ligand. *Inorganica Chim. Acta.* 2012, 382, 72– 78. DOI: 10.1016/j.ica.2011.10.018;
- [13] (d) Colombo, A.; Locatelli, D.; Roberto, D.; Tessore, F.; Ugo Uranyl-Organic One- and Two-Dimensional Assemblies with 2,2'-bipyridine-3,3'-dicarboxylic, biphenyl-3,3',4,4'-tetracarboxylic and bicyclo[2.2.2]oct-7-ene-2,3,5,6-tetracarboxylic Acids. *Cryst. Eng. Comm.* 2012, 14, 131– 137. DOI: 10.1039/c1ce05772c.
- [14] Bourgogne, C.; Massona, P.; Nicouda, F.; Brasselet, S.; Zyss, J., Investigation of New Push-Pull Pyridine--Oxide Derivatives as 1D NLO Chromophores with Vanishing Dipole Moment. *Synth. Met.* 2001, 124, 213– 216. DOI: 10.1016/s0379-6779(01)00453-2.
- [15] Pugachev, M. V.; Pavelyev, R. S.; Nguyen, T. N. T.; Iksanova, A. G.; Lodochnikova, O. A.; Shtyrin, Yu. G., Synthesis and Antitumor Activity of Pyridoxine Monoalkenyl Derivatives. *Russ. Chem. Bull.* 2016, 65, 532– 536. DOI: 10.1007/s11172-016-1333-z.
- [16] (a) Marder, S. R.; Perry, J. W.; Schaefer, W. P., Synthesis of Organic Salts with Large Second-Order Optical Nonlinearities. *Science* 1989, 245, 626– 628. DOI: 10.1126/science.245.4918.626;
- [17] (b) Cole, J. M.; Edwards, T.-C.; Lin, A. J.; Piltz, R. O.; Depotter, G.; Clays, K.; Lee, S.-C.; Kwon, O.-P., Concerted Mitigation of O...H and C(π)...H Interactions Prospects Sixfold Gain in Optical Nonlinearity of Ionic Stilbazolium Derivatives. *Appl. Mater. Interfaces.* 2015, 7, 4693– 4698. DOI: 10.1021/am508213c.
- [18] Kleinpeter, E.; Bölke, U.; Kreicberga, E., Quantification of the Pushe Pull Character of Azodyes and a Basis for Their Valuation as Potential Nonlinear Optical Materials. *Tetrahedron* 2010, 66, 4503– 4509. DOI: 10.1016/j.tet.2010.04.067.
- [19] (a) Shtyrin, Y. G.; Petukhov, A. S.; Strel'nik, A. D.; Garipov, M. R.; Lodochnikova, O. A.; Litvinov, I. A.; Morozov, O. A.; Lovchev, A. V., Derivatives of Pyridoxine with Nonlinear Optical Properties. *RU Patent*, 2501801, 2012;
- [20] (b) Strel'nik, A. D.; Garipov, M. R.; Petukhov, A. S.; Shtyrin, N. V.; Lodochnikova, O. A.; Litvinov, I. A.; Naumov, A. K.; Morozov, O. A.; Klimovitskii, A. E.; Shtyrin, Y. G., Structural and Spectral Characterization of Novel Non-centrosymmetric 2,4-dinitrobenzene Derivative. *Spectrochim. Acta - Part A Mol. Biomol. Spectrosc.* 2014, 117, 793– 797.
- [21] Ziganshin, A. U.; Kalinina, O. S.; Strel'nik, A. D.; Garipov, M. R.; Koshkin, S. A.; Ziganshina, L. E.; Shtyrin, Y. G., Antagonistic Properties of New Non-Phosphorylated Derivatives of Nitrogen-Containing Heterocycles Towards P2 Receptors. *Int. J. Pharmacol.* 2015, 11, 400– 404. DOI: 10.3923/ijp.2015.400.404.
- [22] Korytnyk, W., A Seven-Membered Cyclic Ketal of Pyridoxol. *J. Org. Chem.* 1962, 27, 3724– 3726. DOI: 10.1021/jo01057a532.
- [23] (a) Shtyrin, N. V.; Strel'nik, A. D.; Sysoeva, L. P.; Lodochnikova, O. A.; Klimovitskii, E. N.; Shtyrin, Y. G., New Synthetic Method for 2,3,4-tris(hydroxymethyl)- 6-methylpyridin-5-ol. *Russ. J. Org. Chem.* 2009, 45, 1266– 1268. DOI: 10.1002/chin.201009147;
- [24] (b) Shtyrin, N. V.; Lodochnikova, O. A.; Pugachev, M. V.; Madzhidov, T. I.; Sysoeva, L. P.; Litvinov, I. A.; Klimovitskii, E. N.; Shtyrin, Y. G., Theoretical and Experimental Study on Cyclic 6-methyl-2,3,4-tris(hydroxymethyl)pyridin-5-ol acetones. *Russ. J. Org. Chem.* 2010, 46, 561– 567. DOI: 10.1134/s1070428010040202.
- [25] Shtyrin, N. V.; Lodochnikova, O. A.; Shtyrin, Y. G., Regioisomeric Oximes and Thiosemicarbazones Derived from 6-substituted Pyridoxines. *Mendeleev Commun.* 2012, 22, 169– 170. DOI: 10.1016/j.mencom.2012.05.021.
- [26] Antonioletti, R.; Bonadies, F.; Ciammaichella, A.; Viglianti, A., Lithium Hydroxide as Base in the Wittig Reaction. A Simple Method for Olefin Synthesis. *Tetrahedron* 2008, 64, 4644– 4648. DOI: 10.1016/j.tet.2008.02.091.

- [27] Kolodiazhnyi, O. I., Phosphorus Ylides: Chemistry and Application in Organic Synthesis; Wiley-VCH: Weinheim, 1999.
- [28] Katritzky, A. R.,; Kucharska, H. Z.,; Tucker, M. J.,; Wuest, H. M., Arylazo Derivatives of Pyridoxine. J. Med. Chem. 1966, 9, 620– 622. DOI: 10.1021/jm00322a046.
- [29] Huang, K.-S.,; Britton, D.,; Etter, M. C.,; Byrnx, S. R., Synthesis, Polymorphic Characterization and Structural Comparisons of the Non-Linear Optically Active and Inactive Forms of Polymorphs of 3-(nitroanilino)cycloalk-2-en-1-ones. J. Mater. Chem. 1996, 6, 123– 129. DOI: 10.1039/jm9960600123.
- [30] Shemla, D.,; Zyss, J., Nonlinear Optical Properties of Organic Molecules and Crystals; Academic Press: London, 1987.
- [31] Shtyrlin, Y. G.,; Petukhov, A. S.,; Strel'nik, A. D.,; Nikitina, E. V.,; Garipov, M. R., Antibacterial Compounds Based on Sulfanilic acid and Pyridoxine. RU Patent, 2480471, 2013.
- [32] Morozov, O. A.,; Naumov, A. K.,; Lovchev, A. V.,; Garipov, M. R., Analysis of Nonlinear Optical Materials Properties by Simple Powder Technique. J. Phys. Conf. Ser. 2015, 594, 1– 7. DOI: 10.1088/1742-6596/594/1/012037.